

QUANTUM $SU(2)$ -INVARIANTS FOR THREE-MANIFOLDS ASSOCIATED WITH NON-TRIVIAL COHOMOLOGY CLASSES MODULO TWO

HITOSHI MURAKAMI

ABSTRACT. We show an integrality of the quantum $SU(2)$ -invariant associated with a non-trivial first cohomology class modulo two.

For a given Lie group G and an integer (level) k , E. Witten gave an ‘invariant’ of three-manifolds by using the path integral [33]. Mathematically rigorous proof of its existence was given by several people. See for example [26, 7, 4, 32, 29]. There are also some refinements for these invariants corresponding to various structures of the three-manifolds, such as cohomology classes and spin structures [30, 7, 9, 18, 3, 2].

Some topological properties of the invariants were given especially for $G = SU(2)$. The first striking result was found by R. Kirby and P. Melvin stating that the quantum $SU(2)$ -invariant with level two splits into the sum of the invariants associated with spin structures and each summand can be described in terms of the μ -invariant. They and X. Zhang obtained a topological interpretation for the quantum $SU(2)$ -invariant with level four [8]. In this case the invariant splits into the sum of the invariants associated with the first cohomology classes modulo two and each summand can be described in terms of the Witt invariant and the cohomologies of both the three-manifold and its double cover defined by the cohomology class. Similarly the author gave a topological interpretation for the quantum $SU(3)$ -invariant with level three [18]. It also decomposes into the sum of the invariants associated with the first cohomology classes modulo three and each summand can be described in terms of the cohomologies of the manifold and its triple cover.

If $G = SU(n)$ and n and the level k are coprime, then the invariant splits into the product of the $U(1)$ -invariant [21, 5] and $PSU(n)$ -invariant [7, 10]. If $n + k$ is odd prime and the three-manifold is a rational homology sphere, the invariant is a cyclotomic integer [17, 19, 15, 16, 28]. Similar integrality holds for a three-manifold with positive first Betti number. Moreover one can define the perturbative invariant from the $PSU(n)$ -invariant for rational homology three-spheres [22, 24, 12]. Recently it is proved by T.T.Q. Le that every coefficient of the perturbative invariant for integral homology three-spheres is of finite type in T. Ohtsuki’s sense [23]. (For $n = 2$ case, it is proved in by A. Kriker and B. Spence [11]. In particular its first non-trivial coefficient coincides with (a multiple of) the Casson invariant [1]. (For rational homology three-spheres, Le showed in [12] that this coincides with a

Date: February 1, 2008.

Partially supported by Waseda University Grant for Special Research Projects (No. 98A-623) and Grant-in-Aid for Scientific Research (C) (No. 09640135), the Ministry of Education, Science, Sports and Culture.

multiple of the Casson-Walker invariant [31].) See also [25] for recent developments of these invariants; especially their relation with the LMO invariant [13].

In this paper we consider the quantum $SU(2)$ -invariant with level divisible by four. By V. Turaev, and Kirby and Melvin [7, 30] it decomposes into the sum of the invariants associated with the cohomology classes modulo two. We will consider the case where $(n+2)/2$ is an odd prime.

Let p be an odd prime and M a $\mathbf{Z}/p\mathbf{Z}$ -homology three-sphere. For a cohomology class $\theta \in H^1(M; \mathbf{Z}/2\mathbf{Z})$, let $\tau_{2p}^{SU(2)}(M, \theta)$ be the quantum $SU(2)$ -invariant with level $2p-2$ associated with the $2p$ th root of unity $q = \exp(\pi\sqrt{-1}/p)$. In the previous paper [20] the author proved

Theorem 1. $\tau_{2p}^{SU(2)}(M, 0) \in \mathbf{Z}[1/2, \xi]$ and the ‘coefficient’ of $(\xi-1)$ is congruent to the Casson-Walker invariant modulo p , where $\xi = \exp(2\pi\sqrt{-1}/p)$.

In this paper we will show a weaker result for non-trivial cohomology classes. We will show

Theorem 2.

$$\tau_{2p}^{SU(2)}(M, \theta) \in \begin{cases} (\xi-1)\mathbf{Z}[1/2, \xi] & \text{if } \theta \cup \theta \cup \theta = 0, \\ \sqrt{-1}(\xi-1)\mathbf{Z}[1/2, \xi] & \text{if } \theta \cup \theta \cup \theta \neq 0. \end{cases}$$

Acknowledgment. The author thanks the organizers, especially Sofia Lambropoulou, of the conference ‘Knots in Hellas’ for their hospitality.

1. PROOF

We put $p = 4u + \varepsilon$ with $\varepsilon = \pm 1$. Put $\xi = \exp(2\pi\sqrt{-1}/p)$, $q = \exp(\pi\sqrt{-1}/p)$ and $s = \exp(\pi\sqrt{-1}/2p)$. Therefore $\xi = q^2 = s^4$.

Suppose that a closed three-manifold M is given by an algebraically split, framed link $\mathcal{L}$ with m components. We also assume that the framing of $\mathcal{L}$ is $(f_1, f_2, \dots, f_m)$ with $f_i \not\equiv 0 \pmod{p}$. Moreover we assume that $f_i \equiv 0 \pmod{2}$ for $i \leq b$ and $f_i \equiv 1 \pmod{2}$ for $b < i \leq m$. Let $\theta \in H^1(M; \mathbf{Z}/2\mathbf{Z})$ be presented by the sublink $\mathcal{L}_\theta = L_1 \cup L_2 \cup \dots \cup L_c$ ($c \leq b$). We put $f_i = 2g_i$ for $1 \leq i \leq b$.

By [7, (8.32) Theorem], $\tau_{2p}^{SU(2)}(M, \theta) = \alpha(\mathcal{L})\Sigma(\mathcal{L})$ where $\alpha(\mathcal{L})$ and $\Sigma(\mathcal{L})$ is given as follows.

$$\begin{aligned} \alpha(\mathcal{L}) &= \left(\sqrt{\frac{1}{p}} \sin\left(\frac{\pi}{2p}\right) \right)^m \exp\left(-\frac{3(p-1)}{8p} \times 2\pi\sqrt{-1}\right)^{\sigma(\mathcal{L})} \\ \Sigma(\mathcal{L}) &= \sum_{\substack{0 < k_i \leq p \\ \mathcal{L}_{\mathbf{k}} = \mathcal{L}_\theta}} 2^{|T_{\mathbf{k}}|} [\mathbf{k}] J(\mathcal{L}, \mathbf{k}), \end{aligned}$$

where $J(\mathcal{L}, \mathbf{k})$ is the colored Jones polynomial of $\mathcal{L}$ with color $\mathbf{k} = (k_1, k_2, \dots, k_m)$ ($0 < k_i \leq p$), $\mathcal{L}_{\mathbf{k}}$ is the sublink of $\mathcal{L}$ consisting of components with the color k_i even, $|T_{\mathbf{k}}|$ is the number of k_i which are less than p and $\sigma(\mathcal{L})$ is the number of positives minus that of negatives in $\{f_1, f_2, \dots, f_m\}$.

After some calculations we can simplify $\alpha(\mathcal{L})$ and $\Sigma(\mathcal{L})$ as follows.

Lemma 1.1 (Lemma 2.2 in [20]).

$$\alpha(\mathcal{L}) = \frac{(\xi^u + \xi^{-u})^m}{(2G(\xi))^m} (-1)^{u\sigma(\mathcal{L}) + \sigma_-(\mathcal{L})(\varepsilon-1)/2} \xi^{3 \times \bar{8}\varepsilon\sigma(\mathcal{L})},$$

where $G(\xi)$ is a Gaussian sum $\sum_{k=0}^{p-1} \xi^{k^2}$, $\bar{8}$ is the inverse of 8 in $\mathbf{Z}/p\mathbf{Z}$ and $\sigma_-(\mathcal{L})$ is the number of negatives in $\{f_1, f_2, \dots, f_m\}$. Note that $G(\xi)$ is of the form $\gamma(\xi - 1)^{(p-1)/2}$ with γ invertible in $\mathbf{Z}[\xi]$.

Lemma 1.2 (see Lemma 2.3 in [20]).

$$\begin{aligned} \Sigma(\mathcal{L}) &= [2]2^c \sum_{n_1, n_2, \dots, n_m=1}^{(p-1)/2} (-1)^{\sum_i n_i} V(L^{(2n_1-1, 2n_2-1, \dots, 2n_c-1, 2n_{c+1}, 2n_{c+2}, \dots, 2n_m)}) \\ &\times \prod_{l=1}^c C_{n_l}(g_l) \prod_{l=c+1}^m S_{n_l}(f_l) \end{aligned}$$

with

$$S_n(f) = \sum_{k=n}^{p-n-1} (-1)^k [2k+1] \xi^{(k^2+k)f/2} \binom{k+n}{k-n}$$

and

$$C_n(g) = \sum_{k=n}^{(p-1)/2} (-1)^k [2k] s^{(4k^2-1)g} \binom{k+n-1}{k-n},$$

where $L^{(d_1, d_2, \dots, d_m)}$ is the (unframed) link obtained from $\mathcal{L}$ by replacing i th component with d_i parallels respecting the zero-framing. For a link L , $V(L)$ is the Jones polynomial defined by the skein relation $qV(L_+) - q^{-1}V(L_-) = (s - s^{-1})V(L_0)$ for the usual skein triple (L_+, L_-, L_0) and by $V(\text{trivial knot}) = 1$.

Proof. The proof is similar to [20, Lemma 2.3] and so details are omitted. $\square$

To prove Theorem 2, we prepare three lemmas.

Lemma 1.3 (Lemma 2.4 in [20]). $S_n(f) \in \mathbf{Z}[\xi]$ is divisible by $(\xi - 1)^{(p-1)/2-n}$ in $\mathbf{Z}[\xi]$.

Lemma 1.4. $\sqrt{-1}^{g+1} C_n(g) \in \mathbf{Z}[\xi]$ is divisible by $(\xi - 1)^{(p-1)/2-n+1}$ in $\mathbf{Z}[\xi]$.

Proof. First note that $s = \sqrt{-1}^\varepsilon \xi^{-\varepsilon u}$. Therefore an easily calculation shows that

$$C_n(g) = \frac{\sqrt{-1}^{-\varepsilon(g+1)} \xi^{\varepsilon u g}}{\xi^u + \xi^{-u}} \sum_{k=n}^{(p-1)/2} (\xi^{-2\varepsilon u k} - \xi^{2\varepsilon u k}) \xi^{gk^2} \binom{k+n-1}{k-n}.$$

Now the proof follows from [14, Lemma 6.2] (see also [22]). $\square$

The following lemma is a generalization of a result of M. Sakuma [27, Theorems 1 and 2], who proves the case where $c = 0$.

Lemma 1.5. $\sqrt{-1}^{c+1} V(L^{(2n_1-1, 2n_2-1, \dots, 2n_c-1, 2n_{c+1}, 2n_{c+2}, \dots, 2n_m)}) \in \mathbf{Z}[\xi]$ is divisible by $(\xi - 1)^{\sum_{i=1}^m n_i - c}$.

Proof. We will show more generally that for a link $L = L_1 \cup L_2$, $V(L_1 \cup L_2^2)$ is divisible by $(\xi - 1)^{\sharp(L_2)}$, where L_2^2 is the two-parallel of L_2 with respect to the zero-framing and $\sharp(L_2)$ is the number of components in L_2 .

We will use L.H. Kauffman's bracket polynomial [6] $\langle \mathcal{L} \rangle$ for a framed link $\mathcal{L}$ defined by

$$\left\langle \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\rangle = A \left\langle \begin{array}{c} \diagup \quad \diagup \\ \diagdown \quad \diagdown \end{array} \right\rangle \quad \left\langle \begin{array}{c} \diagup \quad \diagup \\ \diagup \quad \diagup \end{array} \right\rangle + A^{-1} \left\langle \begin{array}{c} \diagdown \quad \diagdown \\ \diagdown \quad \diagdown \end{array} \right\rangle$$

and

$$\langle \mathcal{O} \rangle = 1,$$

where $\mathcal{O}$ is the trivial knot with zero-framing. Since $V(L) = (-1)^{\sharp(L)+1} \langle \mathcal{L} \rangle \Big|_{A=s^{-1/2}}$, we only have to prove that $\langle \mathcal{L}_1 \cup \mathcal{L}_2^2 \rangle$ is divisible by $(A^2 + A^{-2})^{\sharp(\mathcal{L}_2)}$ (note that $s + s^{-1} = \sqrt{-1}^\varepsilon (\xi^{-4\varepsilon u} - \xi^{4\varepsilon u})$). Here $\mathcal{L} = \mathcal{L}_1 \cup \mathcal{L}_2^2$ is the zero-framed link obtained from $L = L_1 \cup L_2^2$.

By simple calculations we have

$$\begin{aligned} & \left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle - \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle \\ &= (A^2 - A^{-2})(A^2 + A^{-2}) \left\{ \left\langle \begin{array}{c} \text{Diagram 3} \end{array} \right\rangle - \left\langle \begin{array}{c} \text{Diagram 4} \end{array} \right\rangle \right\}, \end{aligned}$$

where dotted lines indicate the connectivity. By induction on $\sharp(\mathcal{L}_2)$, we see that the divisibility of $\langle \mathcal{L}_1 \cup \mathcal{L}_2^2 \rangle$ does not change by double crossing change between a single strand and a double strand. This shows that the divisibility of $\langle \mathcal{L}_1 \cup \mathcal{L}_2^2 \rangle$ is greater than or equal to that of $\langle \mathcal{L}_1 \rangle \langle \mathcal{L}_2^2 \rangle$ which is greater than or equal to that of $\langle \mathcal{L}_2^2 \rangle$, which is greater than or equal to $\sharp(\mathcal{L}_2)$ by [20, Lemma 4.1]. This completes the proof. $\square$

Proof of Theorem 2. We only prove the case where M is obtained from an algebraically split, framed link since the general case follows from T. Ohtsuki's diagonalizing lemma [24, Lemma 2.4].

Now we assume that $\theta \in H^1(M; \mathbf{Z}/2\mathbf{Z})$ is given as described before. From the lemmas above, we easily see that $\sqrt{-1}^{\sum_{i=1}^c g_i} \alpha(\mathcal{L}) \Sigma(\mathcal{L})$ belongs to $\mathbf{Z}[1/2, \xi]$ and divisible by $(\xi - 1)$ (Note that $\sqrt{-1}[2] \in \mathbf{Z}[\xi]$ and divisible by $(\xi - 1)$ and that we need $1/2$ from $\alpha(\mathcal{L})$).

Now from [8], $\theta \cup \theta \cup \theta \equiv \sum_{i=1}^c g_i \pmod{2}$. Therefore $\tau_{2p}^{SU(2)}(M, \theta) \in \mathbf{Z}[\xi]$ if $\theta \cup \theta \cup \theta \equiv 0 \pmod{2}$ and $\sqrt{-1}\mathbf{Z}[\xi]$ otherwise, completing the proof. $\square$

REFERENCES

1. S. Akbulut and J.D. McCarthy, *Casson's Invariant for Oriented Homology 3-spheres - An Exposition*, Mathematical Notes, vol. 36, Princeton University Press, Princeton, 1990.
2. C. Blanchet, *Hecke algebras, modular categories and 3-manifolds quantum invariants*.
3. ———, *Refined quantum invariants for three-manifolds with structure*, Knot theory (Warsaw, 1995), Polish Acad. Sci., Warsaw, 1998, pp. 11–22.

4. C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Three-manifold invariants derived from the Kauffman bracket*, *Topology* **31** (1992), 685–699.
5. T. Gocho, *The topological invariant of three-manifolds based on the $u(1)$ gauge theory*, *Proc. Japan Acad. Ser. A Math. Sci.* **66** (1990), no. 8, 237–239.
6. L.H. Kauffman, *State models and the Jones polynomial*, *Topology* **26** (1987), 395–407.
7. R. Kirby and P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin–Turaev for $sl(2, \mathbb{C})$* , *Invent. Math.* **105** (1991), 473–545.
8. R. Kirby, P. Melvin, and X. Zhang, *Quantum invariants at the sixth root of unity*, *Comm. Math. Phys.* **151** (1993), 607–617.
9. T. Kohno and T. Takata, *Symmetry of Witten’s 3-manifold invariants for $sl(n, \mathbb{C})$* , *J. Knot Theory Ramifications* **2** (1993), 149–169.
10. ———, *Level-rank duality of Witten’s 3-manifold invariants*, *Progress in algebraic combinatorics* (Fukuoka, 1993), *Math. Soc. Japan, Tokyo*, 1996, pp. 243–264.
11. A. Krieger and B. Spence, *Ohtsuki’s invariants are of finite type*, *J. Knot Theory Ramifications* **6** (1997), no. 4, 583–597.
12. T. T. Q. Le, *On perturbative $PSU(n)$ invariants of rational homology 3-spheres*.
13. T. T. Q. Le, J. Murakami, and T. Ohtsuki, *On a universal perturbative invariant of 3-manifolds*, *Topology* **37** (1998), no. 3, 539–574.
14. X.-S. Lin and Z. Wang, *On Ohtsuki’s invariants of integral homology 3-spheres, I*, *math.QA/9509149*.
15. G. Masbaum and J.D. Roberts, *A simple proof of integrality of quantum invariants at prime roots of unity*, *Math. Proc. Cambridge Philos. Soc.* **121** (1997), 443–454.
16. G. Masbaum and H. Wenzl, *Integral modular categories and integrality of quantum invariants at roots of unity of prime order*, to appear in *J. Reine Angew. Math.*
17. H. Murakami, *Quantum $SU(2)$ -invariants dominate Casson’s $SU(2)$ -invariant*, *Math. Proc. Cambridge Philos. Soc.* **115** (1993), 253–281.
18. ———, *Quantum invariants for 3-manifolds*, *The 3rd Korea-Japan School of Knots and Links* (K.H. Ko and G.T. Jin, eds.), *Proc. Appl. Math. Workshops*, no. 4, 1994, pp. 129–143.
19. ———, *Quantum $SO(3)$ -invariants dominate the $SU(2)$ -invariant of Casson and Walker*, *Math. Proc. Cambridge Philos. Soc.* **117** (1995), 237–249.
20. ———, *Quantum $SU(2)$ -invariants of three-manifolds associated with the trivial first cohomology class modulo two*, preprint, Waseda Univ., 1998.
21. H. Murakami, T. Ohtsuki, and M. Okada, *Invariants of three-manifolds derived from linking matrices of framed links*, *Osaka J. Math.* **29** (1992), 545–572.
22. T. Ohtsuki, *A polynomial invariant of integral homology 3-spheres*, *Math. Proc. Cambridge Philos. Soc.* **117** (1995), no. 1, 83–112.
23. ———, *Finite type invariants of integral homology 3-spheres*, *J. Knot Theory Ramifications* **5** (1996), no. 1, 101–115.
24. ———, *A polynomial invariant of rational homology 3-spheres*, *Invent. Math.* **123** (1996), no. 2, 241–257.
25. ———, *A filtration of the set of integral homology 3-spheres*, *Proceedings of the International Congress of Mathematicians, Vol. II* (Berlin, 1998), 1998, pp. 473–482.
26. N.Yu. Reshetikhin and V.G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, *Invent. Math.* **103** (1991), 547–597.
27. M. Sakuma, *An evaluation of the Jones polynomial of a parallel link*, *Math. Proc. Cambridge Philos. Soc.* **104** (1988), 105–113.
28. T. Takata and Y. Yokota, *The $PSU(N)$ invariants of 3-manifolds are polynomials*, preprint, Kyushu Univ., 1996.
29. V. Turaev and H. Wenzl, *Quantum invariants of 3-manifolds associated with classical simple Lie algebras*, *Internat. J. Math.* **4** (1993), 323–358.
30. V.G. Turaev, *State sum models in low-dimensional topology*, *Proceedings of the International Congress of Mathematicians, (Kyoto, 1990), Vol. I*, *Math. Soc. Japan, Tokyo*, 1991, pp. 689–698.
31. K. Walker, *An Extension of Casson’s Invariant*, *Annals of Mathematics Studies*, vol. 126, Princeton University Press, Princeton, 1992.
32. H. Wenzl, *Braids and invariants of 3-manifolds*, *Invent. Math.* **114** (1993), 235–275.
33. E. Witten, *Quantum field theory and the Jones polynomial*, *Comm. Math. Phys.* **121** (1989), 351–399.

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY,
OHKUBO, SHINJUKU, TOKYO 169-8555, JAPAN, AND MITTAG-LEFFLER INSTITUTE, AURAVÄGEN
17, S-182 62, DJURSHOLM, SWEDEN

E-mail address: hitoshi@uguisu.co.jp